

# Dust during the Reionization

Erik Elfgren<sup>1</sup> and François-Xavier Désert<sup>2</sup>

<sup>1</sup> Department of Physics, Luleå University of Technology, SE-971 87 Luleå, Sweden

<sup>2</sup> LAOG...


Received ;date; / Accepted ;date;

**Abstract.** The possibility that population III stars have reionized the Universe at redshifts greater than 6 has recently gained momentum with WMAP polarization results. Here we analyse the role of early dust produced by these stars and ejected into the intergalactic medium. We show that this dust, heated by the UV and visible background, produces a submillimetre excess with an electromagnetic spectrum compatible with FIRAS cosmic far infrared background. This spectrum, a Doppler spectrum times the  $\nu^2$  dust emissivity law, is generic and does not depend on other detailed dust properties. Arcminute-scale anisotropies could be detected by future experiments such as Planck HFI.

**Key words.** Dust – CMB – Reionization

## 1. Introduction

More accurate measurements of the cosmic microwave background (CMB) implies a need for a better understanding of the different foregrounds. We study the impact of dust in the very early universe  $5 < z < 15$ . WMAP data on the CMB polarization, Kogut et al. (2003) provides a strong evidence for a rather large Thomson opacity during the reionization of the Universe:  $\tau_e = 0.17 \pm 0.04$  (68%). Although the mechanism of producing such an opacity is not fully understood, Cen, R. (2002, 2003) has shown that early, massive population-III (Pop III) stars could ionize the Universe within  $5 < z < 15$  (see figures 1 and 2). Adopting this hypothesis, we discuss the role and observability of the dust that is produced by the Pop III stars. As we can only conjecture about the physical properties and the abundance of this early dust, we adopt a simple dust grain model with parameters deduced from the Milky Way situation. The dust production is simply linked to the ionizing photon production by the stars through their thermal nuclear reactions. The low potential well of the small pre-galactic halos allows the ejected dust to be widely spread in the intergalactic medium. The ionizing and visible photons from the same Pop III stars heat this dust. There are no direct measurements of this dust, but by means of other results the amount of dust can be estimated. A similar study has been done for a later epoch of the universe, in which data are more readily available, Pei et al. (1999). We use a cosmology with  $\Omega_{tot} = \Omega_m + \Omega_\Lambda = 1$ , where  $\Omega_m = \Omega_b + \Omega_{DM} = 0.022/h^2$ ,  $\Omega_b = 0.0226/h^2$  and  $h = 0.72$  as advocated by WMAP, Spergel et al. (2003), using WMAP data in combination with large scale structure observations (2dF-




**Fig. 1.** Total number of ionizing photons produced from Pop III stars per baryon, cf. (Cen, R. 2002, figure 14). The dotted line represents a simplified model with a constant photon production, 8 per  $z$  per baryon. The results are similar.

GRS + Lyman  $\alpha$ ). Furthermore, since  $z \gg 1$  the universe is matter-dominated.

We now proceed to compute the abundance and the temperature of this dust. Consequences on the CMB distortions are then to be discussed.

## 2. Dust Model

Here we assume the dust properties to be similar to what is observed in our galaxy. For simplicity, we suppose spherical dust grains with radius  $a = 0.1 \mu\text{m}$  and density  $\rho_g = 2.3 \cdot 10^3$



**Fig. 2.** Production rate of ionizing photons from Pop III stars per baryon,  $\frac{dn_\gamma}{dz}$ . The odd form between each integer  $z$  is not physical but due to the discretization.

$\text{kg/m}^3$ . The absorption cross section,  $\sigma_\nu$ , between photons and dust can be written as

$$\sigma_\nu = Q_\nu \pi a^2, \quad (1)$$

where we parametrize the frequency dependency as

$$Q_\nu = \begin{cases} Q_0 \frac{a}{a_r} \left(\frac{\nu}{\nu_r}\right)^\beta & \text{submm and infra red (IR),} \\ 1 & \text{visible and ultra violet (UV),} \end{cases} \quad (2)$$

where  $\nu_r$ ,  $a_r$  and  $Q_0$  are normalization constants. There is only one independent constant which means that we can fix  $a_r = 0.1 \mu\text{m}$  and  $\nu_r = c/(40 \mu\text{m})$ . Above the characteristic frequency  $\nu_r$  the spectral index  $\beta = 1$ , below  $\nu_r$   $\beta = 2$ . The exact position of  $\nu_r$  is unimportant for our study because it is above the interesting frequency region.

In the submm and far IR range, the spectral index is constant, and with  $Q_0 = 0.0088$  the assumed opacity agrees well with measurements by FIRAS on cirrus clouds in our galaxy, cf. Boulanger et al. (1996); Fixsen et al. (1998); Lagache et al. (1999). In the visible and UV region, the cross section is independent of the frequency because  $\lambda < 2\pi a$ . In the submm region, the cross section is proportional to the mass of the grain.

In order to evaluate the significance of the dust during the reionization, we calculate the amount of dust present in the universe at a given time. The co-moving relative dust density is  $\Omega_d = \rho_d(z)/((1+z)^3 \rho_c)$ , where  $\rho_d(z)$  is the dust density,  $z$  is the red-shift,  $\rho_c = \frac{3H_0^2}{8\pi G}$  is the critical density ( $H_0$  and  $G$  are Hubble's and Newton's constants, respectively). It evolves as:

$$\frac{d\Omega_d}{dz} = J_+ - J_-, \quad (3)$$

where  $J_+$  and  $J_-$  are the production and the destruction rate respectively.

The Pop III stars produce enough photons for the reionization while burning  $H$  and thus forming metals ( $Li$  and higher). These metals are released in supernovae explosions at the end of the stars short lives ( $\sim 1$  Myr), whereafter they clump together to form dust, Nozawa et al. (2003). Knowing the production rate of ionizing photons to be  $\frac{dn_\gamma}{dz}$  (figure 2), we can

calculate the total photon energy released from the Pop III stars. This can be done by supposing that each photon has an effective energy of  $E_\gamma = c_\gamma \int_{\nu_{ion}}^{\infty} d\nu h\nu B_\nu(T_*) / \int_{\nu_{ion}}^{\infty} d\nu B_\nu(T_*)$ , where  $h\nu_{ion} = 13.6$  eV and  $B_\nu(T_*)$  is the spectrum of a star with temperature  $T_*$ . The energy of the non-ionizing photons is included through  $c_\gamma = u_{tot}/u_{\nu > \nu_{ion}}$  ( $u$  is the energy from the star). A Pop III star has  $T_* \sim 80\,000$  K (Shioya et al. 2002, page 9) which gives  $E_\gamma \approx 36$  eV. Note that for other reasonable star temperatures,  $E_\gamma$  does not vary significantly,  $E_\gamma|_{60 \times 10^3 K} \approx 36$  eV and  $E_\gamma|_{100 \times 10^3 K} \approx 40$  eV. Hence, the total Pop III photon energy production is  $\frac{dn_\gamma}{dz} E_\gamma$  per baryon per  $z$ . For each consumed nucleon, we assume that a nuclear energy of  $E_r = 7$  MeV is released as radiation, which means that the nucleon consumption rate is  $\frac{dn_\gamma}{dz} E_\gamma / E_r$  nucleons per baryon per  $z$ . If  $f_d$  is the fraction of the consumed baryon mass that becomes interstellar dust, (some of the metal atoms will remain with the core, some will be released as gas) the co-moving dust production rate will be

$$J_+ = f_d \Omega_b \frac{dn_\gamma}{dz} E_\gamma / E_r. \quad (4)$$

A dust grain will eventually be destroyed, e.g. by collision, by supernova shockwaves or by cosmic rays, see Draine & Salpeter (1979) for further discussion. If a dust grain has a lifetime of  $\Delta t$  we can write the dust destruction rate as

$$J_- = \frac{\Omega_d(z)}{\Delta t} \frac{dt}{dz} \approx - \frac{\Omega_d(z)}{\Delta t H_0 \Omega_m^{1/2} (z+1)^{5/2}} \quad (5)$$

since the universe is matter dominated for  $5 < z < 15$ .

Solving eq 3 gives the dust density evolution

$$\Omega_d(z) = \int_z^{z_i} J_+(z') \frac{Y(z')}{Y(z)} dz', \quad (6)$$

where  $z_i = 20$  is the beginning of the dust formation (see figure 1) and

$$Y(z) = \exp\left(\frac{3}{2} \frac{(z+1)^{-3/2}}{\Omega_m^{1/2} \Delta t H_0}\right). \quad (7)$$


We note that the source term  $J_+$  is modulated by the destruction term  $\frac{Y(z')}{Y(z)}$ . The dust density is plotted in figure 3 where we note a strong dependency on the dust lifetime. In local dust  $\Delta t \sim 100$  Myr, Draine & Salpeter (1979). However, the uncertainty is rather large, according to Draine (1990),  $\Delta t = 30$  Myr – 10 Gyr, depending on the environment. Note, however, that the density at the reionization red-shifts is much lower than in the interstellar medium in the Milky Way.

## 3. Results and Discussion

### 3.1. Metallicity

If we suppose that most of the metals were ejected as dust (not as gas) the metallicity comes from the dust grains. The metallicity is directly obtained through the produced dust. By letting  $\Delta t \rightarrow \infty$  ( $\Delta t = 10$  Gyr is good enough) we find the metallicity:

$$\frac{Z}{Z_\odot} = \frac{\Omega_d(\Delta t \rightarrow \infty)/\Omega_b}{0.02} \approx 1147 \cdot \Omega_d(\Delta t \rightarrow \infty) \quad (8)$$



**Fig. 3.** The co-moving relative dust density evolution  $\Omega_d = \rho_{dust}/\rho_c$ , for  $f_d = 1$ . The sink at  $z = 6$  and  $9$  for  $\Delta t \leq 0.1$  Gyr is due to the fact that  $\Delta z = 1$  is not a constant time interval.

or in absolute terms  $Z \approx 22.9\Omega_d$ . At  $z = 5$  we have  $\Omega_d = 2.5 \cdot 10^{-5} f_d$ , which gives  $Z \approx 5.7 \cdot 10^{-4} f_d = 0.029 f_d Z_\odot$ .


There are not much data available for  $z > 5$ . Metal poor stars in our galaxy are one point of reference, absorption lines in the Ly $\alpha$  spectrum from quasars are another one. The lowest metallicities found in stars in the Milky Way are  $Z/Z_\odot \sim 0.01$ , Depagne et al. (2002). The Ly $\alpha$  forest suggests (Songaila & Cowie 2002, figure 13) that  $Z/Z_\odot \sim 0.003$  for  $z \sim 4.5$  assuming that  $[\text{Fe}/\text{H}] \approx \log(Z/Z_\odot)$  as suggested by (VandenBerg et al. 2000, page 432). This is a factor 10 lower than our value if  $f_d = 1$ . However, this might be lower than the actual value, cf. (Pettini et al. 1997, figure 4).

In heavy stars, virtually all the helium is consumed, producing metals. For simplicity (and lack of data), we assume that all the ejected metals clump to form dust,  $f_d \approx f_{eject}$ . This means that  $f_d$  will almost entirely depend on the dust ejection rate in the supernova explosion. In Iwamoto et al. (1998) a detected hypernova of mass  $M \sim 14M_\odot$  seems to have  $f_{eject} \gtrsim 0.7$ . Furthermore, according to a dust production model by Nozawa et al. (2003),  $f_d \approx 0.2 - 0.3$ . At the same time, some of the stars will become black holes, not ejecting any metals, Heger & Woosley (2002), decreasing  $f_d$ . Currently this decrease is largely unknown.

In summary, the mass fraction of the produced metals in the Pop III stars, having become interstellar dust, should be  $f_d = 0.1 - 0.3$ . In the following we use the more conservative  $f_d = 0.1$ , in agreement with the Ly $\alpha$  forest measurements, unless otherwise stated.

### 3.2. Dust Opacity

With our model for the dust density evolution, we want to calculate the opacity of the dust, as seen by the CMB. This will tell us how much the CMB spectrum is altered by the passage through the dust.



**Fig. 4.** Opacity  $\tau$  with dust evolution taken into account.

The dust opacity is given by

$$\begin{aligned} \tau_\nu &= c \int dz \frac{dt}{dz} \sigma_{\nu_e} n_d(z) \\ &= \frac{Q_0 c}{\sqrt{\Omega_m} a_r H_0} \frac{3 \rho_c}{4 \rho_g} \int dz \left( \frac{\nu}{\nu_r} \right)^{\beta_{\nu_e}} \Omega_d(z) (z+1)^{1/2+\beta_{\nu_e}}, \end{aligned} \quad (9)$$

where  $\nu$  ( $\nu_e$ ) is the observed (emitted) frequency and  $\nu = \nu_e/(1+z)$ . The dust number density is  $n_d(z) = (1+z)^3 \times \rho_c \Omega_d(z)/m_g$  where  $m_g = \frac{4\pi a^3}{3} \rho_g$  is the grain mass. We note (from  $\tau \propto \Omega_d$ ) that  $\tau$  is proportional to the parameter  $f_d$ .

The resulting opacity can be seen in figure 4. We note that the opacity is small,  $\tau \ll 1$ . The smooth knee is due to the change of  $\beta$  at the redshifted  $\nu_r$ , see section 2, but this is not in the spectral range of the CMB. The differential opacity  $d\tau/dz$  is plotted in figure 5 for  $\lambda = 1$  mm. We see that with a short dust lifetime, the dust density falls off almost immediately (in terms of  $z$ ). However, for longer lifetimes, the early dust could still play a certain role for  $z < 3$ . This could eventually contribute to dimming of distant objects. We also note the impact of the expansion of the universe in decreasing the dust density and thus the opacity.

### 3.3. Dust Temperature

In order to deduce the equilibrium temperature of the dust, we write the balance between the absorbed CMB, the absorbed starlight and the emitted IR light from the dust:


$$P_d = P_* + P_{CMB}. \quad (11)$$

The powers  $P_d$  and  $P_{CMB}$  can be written as

$$P_X = 4\pi \int_0^\infty d\nu_e \sigma_{\nu_e} B_{\nu_e}(T_X), \quad (12)$$

where  $B_\nu$  is a Planck blackbody spectrum and  $X = \{\text{CMB}, d\}$ . In the wave-length range considered, the spectral index  $\beta = 2$ . Equation 11 can now be solved for the dust temperature analytically in the submm range:

$$T_d^{4+\beta} = T_{*eff}^{4+\beta} + T_{CMB}^{4+\beta}, \quad (13)$$



**Fig. 5.** The differential opacity  $d\tau/dz$  at  $\lambda = 1$  mm for different dust lifetimes.

where the effective temperature is defined by

$$T_{*eff}^{4+\beta} = \frac{P_*}{4\pi C_\beta \sigma_\nu (c/\nu)^\beta 2hc^2 (k_B/hc)^{4+\beta}} \quad (14)$$

and the constant  $C_\beta = (\beta + 3)! \sum_{k=1}^{\infty} k^{-(4+\beta)}$ . This means that  $C_0 \approx 6.494$ ,  $C_1 \approx 24.89$  and  $C_2 \approx 122.1$ .

The absorbed power density,  $P_*$  from the radiation of Pop III stars peaks in the UV-region and can be approximated by

$$P_* = \sigma_{UV} u_*(z) c, \quad (15)$$

where  $\sigma_{UV}$  is the dust-photon cross section in the UV region and the energy density is

$$u_*(z) = f_{esc} \int_{z_i}^z dz' \frac{dn_\gamma}{dz'} E_\gamma \frac{1+z}{1+z'}, \quad (16)$$


where  $f_{esc}$  is the escape fraction of photons from the star halos. We neglect the loss of photons due to the reionization itself.  $E_\gamma \frac{1+z}{1+z'}$  is the effective energy of the photon emitted at  $z'$  and then redshifted to  $z$ . According to [Cen, R. \(2003\)](#),  $f_{esc} = 0.3$  gives an electron opacity  $\tau_e \approx 0.13$  which is within one standard deviation of the results by WMAP. Hereafter, we adopt this value of  $f_{esc}$ .

The energy density of the ionizing photons are compared to the CMB in figure 6. We note that the star energy density is much less than the CMB energy density at this epoch, and that the curve resembles the accumulated photons in figure 1. Hence, the dust temperature closely follows the CMB temperature, see figure 7 and equation 12.


### 3.4. Observed Intensity

Now we proceed to compute the average intensity (monopole term) of the submm and microwave background which is made of the CMB and early dust emission. The simple radiative transfer of the CMB through the uniform dust screen yields the following observed intensity:

$$i_\nu = e^{-\tau_\nu} \left[ B_\nu(T_{CMB}) + \int_0^{\tau_\nu} e^{\tau_e} \frac{B_\nu(T_d(z))}{(1+z)^3} d\tau_e \right]. \quad (17)$$



**Fig. 6.** Energy density of ionizing photons compared to  $u_{CMB} = 4\sigma_B T_{CMB}^4/c$ , where  $\sigma_B$  is Stefan-Boltzmann's constant.




**Fig. 7.** The dust temperature is plotted against the CMB temperature with the relative quantity  $(T_d - T_{CMB})/T_{CMB}$ .

From figures 4 and 7, we see that the opacity is small, ( $\tau \ll 1$ ) and the dust temperature is only slightly higher than the CMB temperature ( $T_d \gtrsim T_{CMB}$ ). This gives the following formula for the excess intensity relative to the unperturbed CMB:

$$\begin{aligned} \Delta i_\nu &\equiv i_\nu - B_\nu(T_{CMB}) \\ &= T_{CMB} \frac{dB_\nu}{dT} \Big|_{T=T_{CMB}} \int_0^{\tau_\nu} \frac{T_d(z) - T_{CMB}(z)}{T_{CMB}(z)} d\tau_e, \end{aligned} \quad (18)$$

where  $T_{CMB}$  is the CMB temperature today. The integrand is plotted in figure 7. We note that a new component is added to the primary CMB spectrum. Equation 18 tells us that it has a specific spectrum which is the product of a Doppler effect spectrum (like primary anisotropies) and a  $\nu^2$  power law. This effect is mostly visible in the submm range and has a vanishing contribution in the radio domain.

In figure 8, the excess intensity is plotted along with the extragalactic background measured by FIRAS, [Puget et al. \(1996\)](#); [Fixsen et al. \(1998\)](#); [Lagache et al. \(1999\)](#). Depending



**Fig. 8.** Comparison of the modeled intensity for the early dust emission in excess of the CMB with the observed FIRAS spectrum (dashed red curve) of the cosmic far IR background as detailed by Lagache et al. (1999).

on the dust destruction rate (parametrized by the dust lifetime  $\Delta t$ ), the computed early dust background can be an important part of the observed background from  $400 \mu\text{m}$  up to the mm wave-length. Most of the far IR background can now be explained by a population of  $z = 0$  to  $z = 3$  luminous IR galaxies, Gispert et al. (2000), the submm fraction of this background could arise from larger redshift dust emission as suggested by figure 8.

In order to check our results, we calculate the co-moving luminosity density of the dust in the submm region and compare it with (Gispert et al. 2000, figure 4). We find them compatible.

### 3.5. Discussion

Just like the Thomson scattering during reionization, early dust will also tend to erase the primordial anisotropies in the CMB. However, due to the much smaller dust opacity (compare  $\tau_d(1\text{mm}) \lesssim 10^{-3}$  and  $\tau_e = 0.17$ ), this effect will be negligible.

The early dust will also introduce a new type of secondary anisotropies with a typical size of a dark matter filament. Here, we only estimate the order of magnitude of this effect. If the co-moving size of the dark matter filament is  $L$ , the angular size is  $3 \cdot (L/5 \text{ Mpc})$  arcminutes at  $z = 10$  which corresponds to multipole number  $\ell \sim 4000 \cdot (L/5 \text{ Mpc})$ . Fortunately, this region in  $\ell$ -space does not contain any primordial fluctuations because of the Silk damping. However, there are other foregrounds in the same region, see Aghanim et al. (2000). If we suppose a contrast of 10% in the dust intensity between dark matter filaments and the void, we obtain values of  $\Delta T/T \approx 5 \times 10^{-7}$  (for  $\lambda = 1 \text{ mm}$ ,  $f_d = 0.1$  and  $\Delta t = 1 \text{ Gyr}$ ). These anisotropies, pending more accurate calculations, clearly are in the range of major expected arcminute secondary anisotropies. They could well be detected by Planck.

The results of these calculations depend only very weakly on the precise dust model assumptions. We have also tried a

different (but similar) shape of the ionizing photon production, figure 1, and found that the results do not vary significantly.

Very little is known about the universe during the reionization epoch. Nevertheless, there are several parameters that could be calculated more accurately.

The two most important parameters in the present model are the dust lifetime,  $\Delta t$  and the mass fraction of the produced metals that are ejected as interstellar dust,  $f_d$ . The dust lifetime could be determined more precisely by making 3D simulations of the dust production in combination with structure formation. The simulations would also give the inhomogeneous dust density evolution. The result would be a better estimate of the aforementioned secondary anisotropies caused by the variations in the dust opacity. A more refined dust grain model, using e.g. a distribution of grain sizes would also be more realistic. If the dust is long-lived, it could also have a certain impact on measurements in the optical and UV region. Finally, we note that most of the results are proportional to the dust density and thus to  $f_d$ . To evaluate  $f_d$  more precisely, we need a better understanding of the typical properties of the first generation of stars, see section 3.1, which is currently a much debated.

## 4. Conclusions

We have shown that early dust ( $5 < z < 20$ ) contributes in the submm and IR to the observed spectrum of the CMB.

The dust is heated by the CMB as well as by superheavy PopIII stars. At the same time, the opacity is so low that the contribution will not be detectable with current instruments. Only by the spectral nature can the dust be distinguished from the primordial CMB anisotropies.

This early dust should show up as small-scale anisotropies with a specific submillimetre spectrum.

## References

- Aghanim, N., Balland, C., & Silk, J. 2000, *A&A*, 357, 1
- Boulanger, F., Abergel, A., Bernard, J.-P., et al. 1996, *A&A*, 312, 256
- Cen, R. 2002, *ApJ*, 591, 12
- Cen, R. 2003, *ApJL*, 591, L5
- Depagne, E., Hill, V., Spite, M., et al. 2002, *A&A*, 390, 187
- Draine, B. T. 1990, in *ASP Conf. Ser. 12: The Evolution of the Interstellar Medium*, 193–205
- Draine, B. T. & Salpeter, E. E. 1979, *ApJ*, 231, 438
- Fixsen, D. J., Dwek, E., Mather, J. C., Bennett, C. L., & Shafer, R. A. 1998, *ApJ*, 508, 123
- Gispert, R., Lagache, G., & Puget, J. L. 2000, *A&A*, 360, 1
- Heger, A. & Woosley, S. E. 2002, *ApJ*, 567, 532
- Iwamoto, K., Mazzali, P. A., Nomoto, K., et al. 1998, *Nature*, 395, 672
- Kogut, A., Spergel, D. N., Barnes, C., et al. 2003, *ArXiv Astrophysics e-prints*
- Lagache, G., Abergel, A., Boulanger, F., Désert, F. X., & Puget, J.-L. 1999, *A&A*, 344, 322
- Nozawa, T., Kozasa, T., Umeda, H., Maeda, K., & Nomoto, K. 2003
- Pei, Y. C., Fall, S. M., & Hauser, M. G. 1999, *ApJ*, 522, 604

- Pettini, M., Smith, L. J., King, D. L., & Hunstead, R. W. 1997, ApJ, 486, 665
- Puget, J.-L., Abergel, A., Bernard, J.-P., et al. 1996, A&A, 308, L5+
- Shioya, Y., Taniguchi, Y., Murayama, T., et al. 2002, ApJ, 576, 36
- Songaila, A. & Cowie, L. L. 2002, AJ, 123, 2183
- Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, ArXiv Astrophysics e-prints
- VandenBerg, D. A., Swenson, F. J., Rogers, F. J., Iglesias, C. A., & Alexander, D. R. 2000, ApJ, 532, 430